

SHORT, NON-REFEREED PAPER

## EFFECT OF FIBRE CONTENT AND SELF-TRASHING TRAITS ON *ELDANA SACCHARINA* WALKER STALK BORER DAMAGE IN SUGARCANE AND IMPLICATIONS FOR RESISTANCE BREEDING

NXUMALO PD<sup>1</sup> AND ZHOU MM<sup>1,2,3</sup><sup>1</sup>South African Sugarcane Research Institute, P/Bag X02, Mount Edgecombe, 4300, South Africa<sup>2</sup>University of the Free State, Plant Sciences Department, Breeding Division, PO Box 339, Bloemfontein, 9300, South Africa<sup>3</sup>University of KwaZulu-Natal, Private Bag X01, Scottsville, 3209, South Africa

Phumla.Nxumalo@sugar.org.za Marvellous.Zhou@sugar.org.za

### Abstract

*Eldana saccharina* (eldana) is the most destructive sugarcane stalk borer in South Africa causing losses estimated at R1 billion per annum. Breeding for resistance complements integrated eldana control strategies. Self-trashing is expected to reduce oviposition sites thus exposing eggs and larvae to predators while high fibre content (>14%) is expected to restrict larval feeding. The study aims to determine the effect of self-trashing and fibre on eldana damage and evaluate implications for resistance breeding. Data on number of bored stalks per genotype were collected from a Stage II trial (TSL16) planted at the SASRI Empangeni Research Station. Significant ( $F=1.86$ ,  $P<0.0001$ ) family effects were observed for eldana damage indicating genetic differences among families. Highly significant ( $F=4.05$ ,  $P<0.0001$ ) family effects were observed for fibre, while those of self-trashing were marginally significant ( $F=1.19$ ,  $P=0.1046$ ) indicating that selection for high fibre and high self-trashing families was possible. Simple linear regression showed a highly significant ( $P<0.0001$ ) negative association representing 0.41% and 0.34% decrease in the number of bored stalks with every unit increase in self-trashing and fibre, respectively. Multiple linear regression showed highly significant ( $P<0.0001$ ) 0.31% (self-trashing) and 0.33% (fibre) decreases in numbers of bored stalks, indicating a higher decrease in eldana damage when both high fibre and high self-trashing were present. Results suggest possible genetic independence between fibre and self-trashing, indicating that both traits can be selected simultaneously during breeding. Additive effects between fibre and self-trashing will accelerate breeding and selection for eldana resistance. Further studies will determine the genetic interactions with commercial traits.

**Keywords:** *Eldana saccharina*, resistance, sugarcane, self-trashing, breeding, fibre content

### Introduction

*Eldana saccharina* Walker (Lepidoptera: Pyralidae) (eldana) is an indigenous insect pest of sugarcane in southern Africa, and is commonly known as the African sugarcane stalk/stem borer. Eldana is the most damaging pest of sugarcane in South Africa causing sugar losses estimated at more than R1 billion per annum (Horton *et al.*, 2002; Assefa *et al.*, 2008; Rutherford, 2015). The use of an Integrated Pest Management (IPM) approach combining chemical control, biological control and cultivation of resistant varieties has shown the potential to alleviate the damaging effects of this borer on sugarcane.

Sugarcane varietal resistance recognised in the early 1970's is an important approach to the control of the sugarcane stalk borer (Keeping and Rutherford, 2004). Several plant defence and/or tolerance mechanisms have been identified, viz. high fibre and hard stalk rindness from research conducted on a few commercial varieties (Mahlanza *et al.*, 2014). More recently, a study by Nxumalo and Zhou (2017) showed reduced eldana damage with every unit increase

in self-trashing. The application of these mechanisms, however, remains limited in breeding programmes.

The objectives of this study were to evaluate the combined effect of self-trashing and fibre content on eldana damage and implications on breeding for eldana resistance.

### Materials and Methods

Data were collected from Stage II TSL16 single lines trial planted at Empangeni Research Station (103 masl; 28°45'S, 31°54'E) in 2016. A randomised completed block design with 164 families replicated three times was used. Each family was made up of several genotypes planted as 8 m single row plots. Genotypes within families were not replicated due to large numbers at the early stage of testing. Data were collected in 2017. At crop maturity, 12 stalks from each genotype (single line) were examined and scored for self-trashing and assessed for eldana damage. Self-trashing was scored from 0 to 3 with 0 representing clinging leaves, 1 representing slight self-trashing, 2 representing intermediate self-trashing and 3 representing highly self-trashing. After scoring for self-trashing, these stalks were trashed, examined for eldana entry and exit holes and the number of damaged stalks was recorded for each genotype. The 12 stalks were then bundled together and sent off to the South African Sugarcane Research Institute (SASRI) fertiliser advisory service (FAS) laboratory, where they were analysed for total fibre expressed as a percentage of sample fresh weight (fibre % cane) using standard methods (Shoones-Muir *et al.*, 2009).

Data were analysed for family effects for eldana damage, self-trashing and fibre using the linear mixed model:

$$Y_{ijk} = \mu + R_j + F_k + RF_{jk} + E_{ijkl} \quad \text{Equation 1}$$

where  $Y_{ijk}$  is the  $i$ th self-trashing/fibre score in the  $j$ th replication of the  $k$ th family measured on the  $l$ th genotype,  $\mu$  is other overall mean,  $R_j$  is the random effect of the  $j$ th replication,  $F_k$  is the fixed effect of the  $k$ th family,  $RF_{jk}$  is the random interaction effect of the  $j$ th replication by the  $k$ th family and is the experimental error for family effects,  $E_{ijkl}$  is the residual error. Families' effects were fixed to estimate means as is practised at this stage in sugarcane breeding.

A simple linear regression was used to determine the relation between eldana damage and fibre as well as eldana damage and self-trashing using the equation,

$$Y_i = a + bx_i + e_i \quad \text{Equation 2}$$

where  $Y_i$  was the eldana damage in the  $i$ th genotype,  $a$  was the intercept,  $b$  was the coefficient or slope,  $x_i$  was the fibre or self-trashing score of the  $i$ th genotype and also the independent variable and  $e_i$  was the error.

Multiple linear regression was used to determine the combined effect of self-trashing and fibre on eldana damage using the equation,

$$Y_i = a + b_1x_{1i} + b_2x_{2i} + e_i \quad \text{Equation 3}$$

where  $a$  was the intercept,  $b_1$  was the slope of the effects of self-trashing,  $b_2$  was the slope of the effect of fibre,  $x_{1i}$  was the self-trashing score of the  $i$ th genotype,  $x_{2i}$  was fibre content of the  $i$ th genotype, and  $e_i$  was the error.

### Results and Discussion

Significant differences ( $F=1.86, P<0.0001$ ) were observed among families for eldana damage suggesting that families with low eldana damage could be identified and selected from these breeding populations (Table 1). There were highly significant ( $F=4.05, P<0.0001$ ) family effects for fibre while those of self-trashing were marginally significant ( $F=1.19, P=0.1046$ ), suggesting genetic variability for these traits (Table 1).

**Table 1. The fixed effects for fibre content and self-trashing.**  
**F-value = F distribution statistic,  $P>F$  = probability of obtaining a larger value of the F statistic.**

Fixed effects			
Effect	Eldana damage	Fibre content	Self-trashing
Family	F-value = 1.86 p-value <0.0001	F-value = 4.05 p-value <0.0001	F-value = 1.19 p-value = 0.1046

Simple linear regression showed a significant negative association representing a 0.41% ( $P<0.0001$ ) and 0.34% ( $P<0.0001$ ) decrease in the number of bored stalks with every unit increase in self-trashing and fibre, respectively (Table 2). This suggests the influence of self-trashing and fibre in reducing eldana damage. The number of bored stalks decreased significantly from 31% (clinging leaves) to 28% (intermediate self-trashing) ( $F=12.80, P<0.0001$ ) for self-trashing while those of fibre decreased significantly from 37% (low fibre) to 22% (high fibre) ( $F=1.85, P<0.0001$ ).

**Table 2. The regression coefficients (estimate), their standard error (SE), t distribution statistic (t-value) and probability of obtaining a larger value of the t-statistic ( $Pr>t$ ).**

Linear regression					
Parameter	DF	Estimate	SE	t-value	Pr>t
Intercept	1	3.620	0.054	66.52	<0.0001
Self-trashing	1	-0.408	0.081	-5.02	<0.0001
Intercept	1	8.133	0.352	23.10	<0.0001
Fibre	1	-0.340	0.025	-13.39	<0.0001
Multiple regression					
Intercept	1	8.129	0.351	23.13	<0.0001
Self-trashing	1	-0.331	0.025	-12.98	<0.0001
Fibre	1	-0.309	0.080	-3.87	0.0001

Multiple linear regression showed highly significant ( $P<0.0001$ ) 0.31% (self-trashing) and 0.33% (fibre) decrease in number of bored stalks, indicating a higher decrease in eldana damage when both high fibre and high self-trashing were present (Table 2). This suggests possible genetic independence between fibre and self-trashing which indicates that both traits can be selected simultaneously during breeding. The number of bored stalks decreased significantly from 31% (clinging leaves) to 29% (intermediate self-trashing) ( $F=10.79, P<0.0001$ ) for self-trashing, while for fibre decreased significantly from 36% (low fibre) to 22% (high fibre) ( $F=15.30, P<0.0001$ ).

In conclusion, results indicated that self-trashing and fibre content traits are associated with the reduction in eldana damage and the interaction of both traits may alleviate damage caused by eldana. These traits may offer an additional resistance mechanism to combat eldana in sugarcane; however, further studies are required to validate the relationship between self-trashing and fibre content with eldana damage. Also, studies are required to determine the interaction of these traits, particularly fibre content, with traits of economic significance.

## REFERENCES

- Assefa Y, Conlong DE, van Den Berg J and Le Rü BP (2008). The wider distribution of *Eldana saccharina* (Lepidoptera: Pyralidae) in South Africa and its potential risk to maize production. *Proc S Afr Sug Technol Ass* 81: 290-297.
- Horton PM, Hearne JW, Apaloo J, Conlong DE, Way MJ and Uys P (2002). Investigating strategies for minimising damage cause by the sugarcane pest *Eldana saccharina*. *Agricultural Systems* 74: 271-286.
- Keeping MG and Rutherford RS (2004). Resistance mechanisms of South African sugarcane to the stalk borer *Eldana saccharina* (Lepidoptera: Pyralidae): A review. *Proc S Afr Sug Technol Ass* 78: 307-312.
- Mahlanza T, Rutherford RS, Snyman SJ and Watt MP (2014). *Eldana saccharina* (Lepidoptera: Pyralidae) resistance in sugarcane (*Saccharum* sp.): Effects of *Fusarium* spp., stalk rind, fibre and nitrogen content. *African Entomology* 22: 810-822.
- Nxumalo PD and Zhou M (2017). Effect of self-trashing on *Eldana saccharina* (Lepidoptera: Pyralidae) damage in sugarcane and implication for resistance breeding. *Proc S Afr Sug Technol Ass* 90: 152-155.
- Rutherford RS (2015). An integrated pest management (IPM) approach for the control of the stalk borer *Eldana saccharina* Walker (Lepidoptera: Pyralidae). In: *IPM for Eldana Control*, South African Sugarcane Research Institute, Mount Edgecombe, 80 pp.
- Shoonees-Muir BM, Ronaldson MA, Naidoo G and Schorn PM (2009). SASTA Laboratory Manual including the Official Methods. Published by the *South African Sugar Technologists' Association*, Mount Edgecombe, Durban, South Africa.